



**Miklós Kassai MD, EBSQ colo**  
Consultant General and  
European Colorectal Surgeon

# Patient Information for Consent

## CR05lite Closure of Loop Ileostomy

Expires end of November 2016  
Issued November 2015

### Local information

Private secretary:  
Fiona Clemets  
Tel: 01534 880 030

Private Clinic:  
Little Grove Clinic  
La Rue De Haut  
St Lawrence  
Jersey  
JE3 1JZ

Get more information and references at [www.aboutmyhealth.org](http://www.aboutmyhealth.org)  
Tell us how useful you found this document at [www.patientfeedback.org](http://www.patientfeedback.org)



[www.rcseng.ac.uk](http://www.rcseng.ac.uk)



[www.rcsed.ac.uk](http://www.rcsed.ac.uk)



[www.asgbi.org.uk](http://www.asgbi.org.uk)



[www.pre-op.org](http://www.pre-op.org)



This document will give you information about closure of loop ileostomy. If you have any questions, you should ask your doctor or other relevant health professional.

### What is closure of a loop ileostomy?

A loop ileostomy is a type of stoma and was made with two ends of your small bowel (see figure 1). It was expected that this would be only temporary and that your bowel ends would be put back together. Your bowels will then open in the usual way.



Figure 1  
A loop ileostomy

### What are the benefits of surgery?

You should be able to open your bowels in the normal way and you should no longer have a stoma bag.

### What does the operation involve?

The operation is performed under a general anaesthetic and usually takes about 45 minutes. Your surgeon will make a cut on your skin around the ileostomy. They will free up the loops of small bowel used to make the ileostomy and then join the two ends back together.

### What complications can happen?

#### 1 General complications

- Pain
- Bleeding
- Infection of the surgical site (wound)
- Blood clots
- Chest infection
- Unsightly scarring
- Developing a hernia

- Difficulty passing urine

#### 2 Specific complications

- Anastomotic leak
- Bowel obstruction
- Diarrhoea

### How soon will I recover?

You should be able to go home after three to five days.

You should feel strong enough to return to normal activities within a few weeks of going home.

Regular exercise should help you to return to normal activities as soon as possible. Before you start exercising, ask the healthcare team or your GP for advice.

Most people make a good recovery. You will be followed up as planned after your original operation.

### Summary

Surgery to join the two ends of your bowel back together should allow you to open your bowels in the normal way. You will no longer need a stoma bag.

### Acknowledgements

Author: Mr Jonathan Lund DM FRCS (Gen. Surg.)  
Illustrations: Medical Illustration Copyright ©  
Medical-Artist.com

**This document is intended for information purposes only and should not replace advice that your relevant health professional would give you.**