

Patient Information for Consent

CR08lite Lateral Internal Sphincterotomy

Expires end of November 2016
Issued November 2015

Local information

Private secretary:
Fiona Clemets
Tel: 01534 880 030

Private Clinic:
Little Grove Clinic
La Rue De Haut
St Lawrence
Jersey
JE3 1JZ

Get more information and references at www.aboutmyhealth.org
Tell us how useful you found this document at www.patientfeedback.org



www.rcseng.ac.uk



www.rcsed.ac.uk



www.asgbi.org.uk



www.pre-op.org

This document will give you information about a lateral internal sphincterotomy. If you have any questions, you should ask your GP or other relevant health professional.

What is an anal fissure?

An anal fissure is a tear in the skin around your back passage (see figure 1). It is a common problem that causes severe pain, especially after a bowel movement. It may also cause bleeding. The condition is associated with spasm of the internal anal sphincter. This reduces the blood supply to the area and prevents healing. The treatment is aimed at breaking this cycle to allow healing to take place.

What are the benefits of surgery?

Surgery should help the anal fissure to heal but is usually recommended if the fissure has not healed with non-surgical treatments.

Are there any alternatives to surgery?

There are simple treatments which may help such as laxatives, ointments, injections of Botox, increasing the amount of fibre in your diet and drinking plenty of fluid.

What does the operation involve?

Sphincterotomy simply means dividing the sphincter. The operation is usually performed under a general anaesthetic and takes about 15 minutes.

Your surgeon will make a small cut on the skin near your back passage. They will cut the lower part of the internal sphincter muscle. This will relieve the spasm in the sphincter, allowing a better blood supply to heal the fissure.

What complications can happen?

1 General complications

- Pain
- Bleeding
- Infection of the surgical site (wound)

2 Specific complications

- Involuntarily passing wind or loose faeces
- Difficulty passing urine
- Permanent incontinence from your bowel




Figure 1
The position of an anal fissure

How soon will I recover?

You should be able to go home the same day. The pain from the fissure should improve rapidly. You should be able to return to work after a few days, depending on your type of work. Regular exercise should help you to return to normal activities as soon as possible. Before you start exercising, ask the healthcare team or your GP for advice.

Most people make a full recovery and can return to normal activities.
The fissure can come back.

Summary

An anal fissure is a common condition that causes a lot of pain. At first, it may be treated with ointments or Botox. If this fails, surgery is the best option for a cure.

Acknowledgements

Author: Mr Jonathan Lund DM FRCS (Gen. Surg.)

Illustrations: Medical Illustration Copyright ©

Medical-Artist.com

This document is intended for information purposes only and should not replace advice that your relevant health professional would give you.