



Miklós Kassai MD, EBSQ colo
Consultant General and
European Colorectal Surgeon

Patient Information for Consent

CR10lite Reversal of Hartmann's Procedure

Expires end of November 2016
Issued November 2015

Local information

Private secretary:
Fiona Clemets
Tel: 01534 880 030

Private Clinic:
Little Grove Clinic
La Rue De Haut
St Lawrence
Jersey
JE3 1JZ

Get more information and references at www.aboutmyhealth.org
Tell us how useful you found this document at www.patientfeedback.org



www.rcseng.ac.uk

www.rcsed.ac.uk

www.asgbi.org.uk

www.pre-op.org

This document will give you information about reversal of Hartmann's procedure. If you have any questions, you should ask your GP or other relevant health professional.

What is Hartmann's procedure?

Hartmann's procedure involves removing a piece of large bowel (colon). It is usually performed as an emergency. The two ends of bowel were not joined together because your surgeon decided the risks were too high. Your surgeon made a colostomy (your large bowel opening onto your skin). The other end was closed as a rectal stump (see figure 1). Your surgeon has decided it is safe to join the two ends of your bowel back together.




Figure 1
A Hartmann's procedure

What are the benefits of surgery?

You should be able to open your bowels in the normal way and you should no longer have a stoma bag.

What does the operation involve?

The operation is performed under a general anaesthetic and usually takes about 90 minutes to two hours.

Your surgeon will make a cut through the scar from the first operation and a cut on your skin around the colostomy. Your surgeon will join the two ends of your bowel back together and place your bowel inside your abdominal cavity.

What complications can happen?

1 General complications

- Pain
- Bleeding
- Infection of the surgical site
- Chest infection
- Unsightly scarring
- Developing a hernia
- Blood clots

2 Specific complications

- Anastomotic leak
- Continued bowel paralysis
- Diarrhoea
- Death

How soon will I recover?

It is usual for your bowel to stop working for a few days.

You should be able to go home after 7 to 10 days. It can take up to three months to return to normal activities.

Regular exercise should help you to return to normal activities as soon as possible. Before you start exercising, ask the healthcare team or your GP for advice.

Most people make a good recovery.

Summary

Reversal of Hartmann's procedure will allow you to open your bowels in the normal way. You will no longer need a stoma bag.

Acknowledgements

Author: Mr Jonathan Lund DM FRCS (Gen. Surg.)
Illustrations: Medical Illustration Copyright ©
Medical-Artist.com

This document is intended for information purposes only and should not replace advice that your relevant health professional would give you.