



Miklós Kassai MD, EBSQ colo
Consultant General and
European Colorectal Surgeon

Patient Information for Consent

GS16lite Surgery for Ingrowing Toenail (adult)

Expires end of November 2016
Issued November 2015

Local information

Private secretary:
Fiona Clemets
Tel: 01534 880 030

Private Clinic:
Little Grove Clinic
La Rue De Haut
St Lawrence
Jersey
JE3 1JZ

Get more information and references at www.aboutmyhealth.org
Tell us how useful you found this document at www.patientfeedback.org



The
Information
Standard

Certified member



RCS

ADVANCING SURGICAL STANDARDS



Excellence in short stay surgery



Preoperative
The
Association

www.rcseng.ac.uk

www.bads.co.uk

www.rcsed.ac.uk

www.asgbi.org.uk

www.pre-op.org

This document will give you information about surgery for an ingrowing toenail. If you have any questions, you should ask your GP or other relevant health professional.

What is an ingrowing toenail?

An ingrowing toenail happens when your toenail grows into the skin around it (see figure 1). The skin can get damaged, causing infection and pain. The problem usually happens to your big toe.



Figure 1
An ingrown toenail

The most common cause of an ingrowing toenail is not cutting your toenails properly. The problem often runs in families as a common cause is having toenails that do not grow straight. Wearing shoes that do not have enough width to fit your toes in their natural position can also cause an ingrowing toenail.

What are the benefits of surgery?

You should no longer have infection and pain.

Are there any alternatives to surgery?

Try to cut your toenails straight across and do not leave sharp edges.

Wear shoes that comfortably fit the shape of your feet.

Your GP or podiatrist may be able to push the skin away from the ingrowing toenail or cut away the piece of toenail that has grown into the skin.

What does the operation involve?

The operation is usually performed under a local anaesthetic.

The operation usually takes about 10 minutes and may involve nail avulsion (removing your whole toenail), wedge excision (removing part of your toenail), nail-bed ablation (removing part of or the whole toenail and then applying a chemical or an electric current to the area of tissue that your toenail grows from), and Zadek's procedure (removing your whole toenail and then cutting away the area of tissue that your toenail grows from).

What complications can happen?

1 General complications

- Pain
- Bleeding
- Unsightly scarring

2 Specific complications

- Infection of the surgical site (wound)
- Infection in the underlying bone

How soon will I recover?

After a short while you will be able to go home. Spend most of the time during the first few days with your leg raised so that the swelling settles. After that, you can usually start to be a little more active.

Do not play sports or do strenuous exercise for two weeks. Keep the dressing dry.

Try to cut your toenails straight across and do not leave sharp edges. Wear shoes that fit the shape of your feet comfortably.

An ingrowing toenail can happen again.

Summary

Surgery to remove an ingrowing toenail should take away infection and pain.

Acknowledgements

Author: Prof Simon Parsons DM FRCS (Gen. Surg.) and Maria Laye, Surgical Care Practitioner
Illustrations: Medical Illustration Copyright © Medical-Artist.com

This document is intended for information purposes only and should not replace advice that your relevant health professional would give you.